

DR. PETER F. REBEIRO (Orcid ID : 0000-0003-1951-9104)

Article type : Viewpoint

Implications of COVID-19 for HIV Research: data sources, indicators, and longitudinal analyses

Peter F. Rebeiro^{1§}, Stephany N. Duda², Kara K. Wools-Kaloustian³, Denis Nash⁴, Keri N. Althoff⁵, on behalf of the International epidemiology Databases to Evaluate AIDS (IeDEA)

- 1 Department of Medicine (Divisions of Infectious Diseases & Epidemiology) & Department of Biostatistics, Vanderbilt University School of Medicine, Nashville, TN, USA
- 2 Department of Biomedical Informatics, Vanderbilt University School of Medicine, Nashville, TN, USA
- 3 Division of Infectious Diseases, Department of Medicine, Indiana University School of Medicine, Indianapolis, IN, USA
- 4 Institute for Implementation Science in Population Health, City University of New York, New York, NY, USA
- 5 Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA

This is the author's manuscript of the article published in final edited form as:

Rebeiro, P. F., Duda, S. N., Wools-Kaloustian, K. K., Nash, D., Althoff, K. N., & International epidemiology Databases to Evaluate AIDS (IeDEA). (2020). Implications of COVID-19 for HIV Research: data sources, indicators and longitudinal analyses. *Journal of the International AIDS Society*, 23(10), e25627. <https://doi.org/10.1002/jia2.25627>

§ Corresponding author:

Peter F. Rebeiro, PhD, MHS

1161 21st Ave. S., A-2200 MCN

Nashville, TN 37232, USA

Phone: +1-615-343-8351

Fax: +1-615-343-6160

Email: p.rebeiro@vanderbilt.edu

E-mail addresses of authors:

PFR: p.rebeiro@vanderbilt.edu

SND: stephany.duda@vanderbilt.edu

KKW: kwools@iu.edu

DN: Denis.Nash@sph.cuny.edu

KNA: kalthoff@jhu.edu

Keywords: HIV care continuum, HIV epidemiology, COVID-19, research design, data sources, longitudinal studies

Word count:

Main text: 962/1000

1 Main text

2

3 *Introduction*

4 Observational research is critical to inform guidelines, policy, and the practice of HIV service
5 delivery.¹ The COVID-19 pandemic has profoundly affected healthcare systems and health behaviors
6 world-wide, including at clinics and research sites that undergird global observational HIV research.²
7 We consider the impact of the COVID-19 pandemic on the capture of relevant HIV data, indicator
8 fidelity, and analytic approaches when investigating effects of COVID-19 itself or accounting for
9 COVID-related changes in service delivery and care-seeking.

10

11 *Data Sources*

12 Observational HIV research relies on robust data sources that accurately reflect the delivery of routine
13 patient care, which is the underlying data-generating mechanism. Due to the COVID-19 pandemic,
14 patient health behaviors and HIV clinical care models have changed. Patients may be unable to attend
15 clinic due to COVID-19 health concerns, reduced transportation, and stay-at-home orders. Clinics
16 globally have responded with increased remote interactions through telehealth, electronic patient
17 portals, social media platforms, and text and email messaging, as well as decentralized antiretroviral
18 treatment (ART) delivery.

19

20 However, new types of care and medication delivery may not be recorded consistently in paper or
21 electronic health record systems and many providers working off-site may not have access to those
22 systems for documentation.^{2,3} Even if these interactions are recorded, data may be inaccessible to
23 researchers if stored in new systems or data fields.⁴ Linked data sources such as pharmacy systems
24 may also experience changes in data quality and content, as prescriptions are dispensed in batches for
25 community delivery or transferred to pharmacies offering reduced-contact dispensing.^{2,3} Mortality
26 and other registries may experience data entry delays due to reporting delays and reduced staffing.
27 These pandemic-related changes are likely to be as heterogeneous across the globe as the pandemic
28 itself, disrupting the data sources researchers have used to assess trends in key HIV-related outcomes,
29 resulting in unreliable and invalid measures of care.²

30

31 *Indicators & Measurement*

32 The HIV care continuum has become the preferred framework for understanding individual
33 movement through various stages of HIV care, from testing and linkage to care, to retention in care,
34 ART receipt, and ultimately, viral suppression. Barriers at various stages of the continuum have been
35 conceptualized as “leaks,” with gaps, delays, and transitions out of care seen as undesirable events
36 that should be mitigated through intervention.^{5,6}

37

38 HIV testing, diagnosis, and linkage to care have been delayed due to suspension or limitation of
39 testing programs during mandatory public quarantine or social distancing measures. Outpatient clinic
40 visits have been shifted to remote encounters when possible, and non-urgent care has been
41 postponed.^{2,7} Individuals that appear to be lost to follow-up at their usual site of care may be seeking
42 care elsewhere and medical records may not travel with them. Delays in ART initiation and refills
43 have occurred due to loss of insurance, limited pharmacy dispensing capabilities, and/or limited
44 outpatient activities. CD4 and viral load monitoring – central to HIV care – may also be delayed due
45 to less available phlebotomy services or co-opting of equipment for COVID-related testing.³

46

47 Both improved data capture and alteration of care continuum metrics may therefore be required to
48 assess the extent of care/service disruptions and reduce measurement error and misclassification.⁸
49 More sensitive definitions of engagement and retention accounting for non-traditional interactions
50 may also be warranted. Many current retention metrics require clinical interactions every 3-6 months,
51 but individuals successfully managed on ART and virally suppressed may need less frequent visits.^{6,9}
52 A conditional retention measure, based on ART receipt and viral suppression prior to clinic service
53 disruption or upon return to clinic, may therefore be more informative about the care continuum than
54 attended or even missed visit counts alone. For example, we could redefine retention status such that
55 an individual would be successfully retained if they were virally suppressed and receiving ART both
56 before local social isolation measures were imposed and after return to the clinic. Such a measure
57 would be a more meaningful indication of the current state of a patient's engagement in the HIV care

continuum, even if their recorded HIV care visits were not frequent enough to meet current retention definitions.

60

Analytic Considerations

In addition to changes in data collection and measurement during study design and conduct, we will also need to use analytic approaches that address the potential for artifactual temporal changes in HIV indicators due to COVID-19, selection biases, and measurement errors in the data-generating mechanisms of the care continuum. If patients attending telehealth visits are not representative of the entire cohort, if outcomes are unreliably ascertained among those lost to care, if certain measures are self-reported remotely instead of being collected on-site, or if discontinuities such as disruptions in care persist, appropriate epidemiologic and biostatistical methods such as inverse probability weighting, multiple imputation, double-sampling, and regression calibration should be considered.^{10–12} To facilitate longer-term trend assessments which span the COVID-19 pandemic, analyses should also accommodate maximum flexibility, for example, through the use of restricted cubic splines, piecewise regression, or parametric mixture models.^{13,14} We must continue to assess the local clinical context to obtain more information relevant to HIV care changes induced by the COVID-19 pandemic and inform these approaches.¹⁵

75

Conclusion

Future HIV-related studies and public health goals require a new COVID-19-informed paradigm for the collection and use of observational cohort data. HIV cohorts must capture pandemic-driven changes in data sources, clinic activities, and local policies to inform analyses. Our ability to leverage epidemiologic evidence to inform clinical, programmatic, and public health practice is only as strong as the inferences derived from these analyses are valid and robust to the challenges in HIV care and research that we now face due to the pandemic. Healthcare organizations and public health agencies should revise HIV care continuum measures and analytic strategies. Funding for such work is critical, even in times of economic crisis, so that COVID-19 does not derail the global fight to End the HIV Epidemic.

86 **Competing interests**

87 The authors declare no competing interests.

88

89 **Authors' contributions**

90 PFR, SND, and KNA developed the idea and wrote the initial draft. KKW and DN revised the
91 manuscript and provided expert input. All authors approved the final manuscript.

92

93 **Acknowledgements**

94 Funding: The International Epidemiology Databases to Evaluate AIDS (IeDEA) is supported by the
95 U.S. National Institutes of Health's National Institute of Allergy and Infectious Diseases (NIAID), the
96 *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, the National
97 Cancer Institute, the National Institute of Mental Health, the National Institute on Drug Abuse, the
98 National Heart, Lung, and Blood Institute, the National Institute on Alcohol Abuse and Alcoholism,
99 the National Institute of Diabetes and Digestive and Kidney
100 Diseases, the Fogarty International Center, and the National Library of Medicine: CCASAnet,
101 U01AI069923; Central Africa, U01AI096299; East Africa, U01AI069911; NA-ACCORD,
102 U01AI069918. Informatics resources are supported by the Harmonist project, R24AI124872. PFR
103 received additional funding from NIAID (K01-AI131895, R21-AI145686).

104

105 Disclaimer: This work is solely the responsibility of the authors and does not necessarily
106 represent the official views of any of the institutions mentioned above.

107

108

109 References

- 110 1. Ford N, Penazzato M, Vitoria M, Doherty M, Davies M-A, Zaniewski E, Tymejczyk O, Egger
111 M, Nash D. The contribution of observational studies in supporting the WHO ‘treat all’
112 recommendation for HIV/AIDS. *J Virus Erad.* 4(Suppl 2):5–8. PMCID: PMC6248853
- 113 2. World Health Organization. Disruption in HIV, Hepatitis and STI services due to COVID-19
114 [Internet]. [cited 2020 Jul 17]. Available from: [https://www.who.int/docs/default-source/hiv-](https://www.who.int/docs/default-source/hiv-hq/presentation-disruption-in-services-international-aids-conference-2020.pdf?sfvrsn=d4bflf87_7)
115 [hq/presentation-disruption-in-services-international-aids-conference-](https://www.who.int/docs/default-source/hiv-hq/presentation-disruption-in-services-international-aids-conference-2020.pdf?sfvrsn=d4bflf87_7)
116 [2020.pdf?sfvrsn=d4bflf87_7](https://www.who.int/docs/default-source/hiv-hq/presentation-disruption-in-services-international-aids-conference-2020.pdf?sfvrsn=d4bflf87_7)
- 117 3. Amimo F, Lambert B, Magit A. What does the COVID-19 pandemic mean for HIV,
118 tuberculosis, and malaria control? *Trop Med Health.* 2020;48:32. PMCID: PMC7218555
- 119 4. Reeves JJ, Hollandsworth HM, Torriani FJ, Taplitz R, Abeles S, Tai-Seale M, Millen M, Clay
120 BJ, Longhurst CA. Rapid Response to COVID-19: Health Informatics Support for Outbreak
121 Management in an Academic Health System. *J Am Med Inform Assoc.* 2020 Mar 24; PMCID:
122 PMC7184393
- 123 5. World Health Organization. Consolidated guidelines on the use of antiretroviral drugs for
124 treating and preventing HIV infection: recommendations for a public health approach.
125 Consolidated guidelines on the use of antiretroviral drugs for treating and preventing HIV
126 infection: recommendations for a public health approach. 2016;(Ed. 2).
- 127 6. Joint United Nations Programme on HIV/AIDS (UNAIDS). 90-90-90: An ambitious treatment
128 target to help end the AIDS epidemic. 2014;40.
- 129 7. Chenneville T, Gabbidon K, Hanson P, Holyfield C. The Impact of COVID-19 on HIV
130 Treatment and Research: A Call to Action. *Int J Environ Res Public Health.* 2020 Jun;17(12).
131 PMCID: PMC7345635

- 132 8. Nash D. Designing and Disseminating Metrics to Support Jurisdictional Efforts to End the
133 Public Health Threat Posed by HIV Epidemics. *Am J Public Health*. 2020 Jan;110(1):53–57.
134 PMCID: PMC6893332
- 135 9. Rebeiro PF, Horberg MA, Gange SJ, Gebo KA, Yehia BR, Brooks JT, Buchacz K, Silverberg
136 MJ, Gill J, Moore RD, Althoff KN, Research NAACC on, Design (NA-ACCORD). Strong
137 agreement of nationally recommended retention measures from the Institute of Medicine and
138 Department of Health and Human Services. *PLoS One*. 2014;9(11):e111772. PMCID:
139 PMC4222946
- 140 10. Cole SR, Hernán MA. Constructing inverse probability weights for marginal structural models.
141 *Am J Epidemiol*. 2008;168(6):656–64. PMCID: PMC2732954
- 142 11. Yiannoutsos CT, An M-WW, Frangakis CE, Musick BS, Braitstein P, Wools-Kaloustian K,
143 Ochieng D, Martin JN, Bacon MC, Ochieng V, Kimaiyo S. Sampling-based approaches to
144 improve estimation of mortality among patient dropouts: experience from a large PEPFAR-
145 funded program in Western Kenya. *PLoS One*. 2008;3(12):e3843. PMCID: PMC2585792
- 146 12. Moscoe E, Bor J, Bärnighausen T. Regression discontinuity designs are underutilized in
147 medicine, epidemiology, and public health: a review of current and best practice. *J Clin*
148 *Epidemiol*. 2015;68(2):122–33.
- 149 13. Shepherd BE, Rebeiro PF, Caribbean SA network for H epidemiology (CCASAnet) Central.
150 Assessing and interpreting the association between continuous covariates and outcomes in
151 observational studies of HIV using splines. *J Acquir Immune Defic Syndr*. 2016;74(3):e60–e63.
152 PMCID: PMC5303133
- 153 14. Lau B, Cole SR, Gange SJ. Parametric mixture models to evaluate and summarize hazard ratios
154 in the presence of competing risks with time-dependent hazards and delayed entry. *Stat Med*.
155 2011;30(6):654–65. PMCID: PMC3069508

- Accepted Article
- 156 15. Duda SN, Farr AM, Lindegren ML, Blevins M, Wester CW, Wools-Kaloustian K, Ekouevi DK,
157 Egger M, Hemingway-Foday J, Cooper DA, Moore RD, McGowan CC, Nash D, International
158 Epidemiologic Databases to Evaluate AIDS (IeDEA) Collaboration. Characteristics and
159 comprehensiveness of adult HIV care and treatment programmes in Asia-Pacific, sub-Saharan
160 Africa and the Americas: results of a site assessment conducted by the International
161 epidemiologic Databases to Evaluate AIDS (IeDEA) Collaboration. J Int AIDS Soc.
162 2014;17:19045. PMCID: PMC4268491